

models used for determining upper bound concentrations of pesticides in surface water and ground water.

The acute drinking water levels of concern (DWLOCs) are 1.3 parts per million (ppm) for the U.S. population, and 0.38 ppm for the most exposed population subgroup, children (1–6 years). The estimated maximum concentration of cymoxanil in surface water (8.15 ppb) derived from GENECC is much lower than the acute DWLOC. Therefore, one can conclude with reasonable certainty that residues of cymoxanil in drinking water will not contribute significantly to the aggregate acute human health risk.

The chronic DWLOCs are 1.4 ppm for the U.S. population and 0.4 ppm for the most sensitive subgroup, children (1–6 years). The DWLOCs are substantially higher than the GENECC 56-day estimated environmental concentration of 0.37 ppb for cymoxanil in surface water. Therefore, one can conclude with reasonable certainty that residues of cymoxanil in drinking water do not contribute significantly to the aggregate chronic human health risk.

**2. Non-dietary exposure.** Cymoxanil products are not labeled for residential non-food uses, thereby eliminating the potential for residential exposure. Non-occupational, non-dietary exposure for cymoxanil has not been estimated because the proposed products are limited to commercial crop production. Therefore, the potential for non-occupational exposure is insignificant.

#### *D. Cumulative Effects*

EPA's consideration of a common mechanism of toxicity is not necessary at this time because there is no indication that toxic effects of cymoxanil should be cumulative with those of any other chemical compounds or with each other. Cymoxanil is a unique cyanoacetamide and is chemically unrelated to any other commercial plant disease control agent. Its biochemical mode of action on fungi appears to be unique; it is theorized to act through inhibition of multiple cellular processes, but a definitive mechanism has not been completely elucidated. Similarly, the mechanism of action underlying observed toxicological effects in mammals is not fully characterized and there is no reliable information to suggest that cymoxanil has a mechanism of toxicity in common with any other compound.

Given the distinct chemical and toxicological profile of cymoxanil, its low acute toxicity, absence of genotoxic, oncogenic, developmental, or reproductive effects, and low exposure potential, the expression of cumulative

human health effects with any other natural or synthetic pesticide is not anticipated.

#### *E. Safety Determination*

**1. U.S. population.** Dietary and occupational exposure will be the major routes of exposure to the U.S. population for cymoxanil, and ample margins of safety have been demonstrated for both.

For cymoxanil, assuming 30% crop treated and residues estimated based on field trial results, the chronic dietary exposure for the overall U.S. population is estimated to be 0.000063 mg/kg/day, using 0.2 percent of the RfD. For acute dietary exposure, the estimated exposure is 0.000475 and 0.001789 at the 99<sup>th</sup> and 99.9<sup>th</sup> percentiles, which will utilize 1.19 and 4.47%, respectively, of the RfD for the overall U.S. population. The ground application margin of exposure (MOE) was 7,814 for mixers/loaders and 1,430 for applicators. The aerial application MOE was 3,907 for mixers/loaders and 38,763 for applicators. The MOE for flaggers was 10,916. Based on the completeness and reliability of the toxicity data and the conservative exposure assessments, there is reasonable certainty that no harm will result from the aggregate exposure of residues of cymoxanil including all anticipated dietary exposure and all other non-occupational exposures.

**2. Infants and children.** Chronic dietary exposure of cymoxanil for the most highly exposed children's subpopulations are: 0.000074 mg/kg/day for children 1–6 years and 0.000068 mg/kg/day for children 7–12 years, representing 0.2% of the chronic reference dose (cRfD) for each subpopulation. Exposure for all infant subpopulations was negligible. For acute dietary exposure of cymoxanil, the %RfD for children 1–6 years is 1.44 at the 99<sup>th</sup> percentile and 5.24 at the 99.9<sup>th</sup> percentile. For non-nursing infants (>1 yr.), the %RfD is 0.46 at the 99<sup>th</sup> percentile and 1.50 at the 99.9<sup>th</sup> percentile. There are no residential uses of cymoxanil; it is extremely unlikely that drinking water will be contaminated. Based on the completeness and reliability of the toxicity data base, the lack of toxicological endpoints of special concern, the lack of any indication that children are more sensitive than adults to cymoxanil, and the conservative exposure assessment, there is a reasonable certainty that no harm will result to infants and children from the aggregate exposure of residues of cymoxanil, including all anticipated dietary exposure and all other non-

occupational exposures. Accordingly, there is no need to apply an additional safety factor for infants and children.

#### *F. International Tolerances*

To date, no international tolerances exist for cymoxanil.

[FR Doc. 03–4257 Filed 2–27–03; 8:45 am]

BILLING CODE 6560–50–S

## FEDERAL EMERGENCY MANAGEMENT AGENCY

[FEMA–1445–DR]

### Alaska; Amendment No. 3 to Notice of a Major Disaster Declaration

**AGENCY:** Federal Emergency Management Agency (FEMA).

**ACTION:** Notice.

**SUMMARY:** This notice amends the notice of a major disaster declaration for the State of Alaska, (FEMA–1445–DR), dated December 4, 2002, and related determinations.

**EFFECTIVE DATE:** February 12, 2003.

#### **FOR FURTHER INFORMATION CONTACT:**

Magda Ruiz, Response and Recovery Directorate, Federal Emergency Management Agency, Washington, DC 20472, (202) 646–2705 or [Magda.Ruiz@fema.gov](mailto:Magda.Ruiz@fema.gov).

**SUPPLEMENTARY INFORMATION:** The notice of a major disaster declaration for the State of Alaska is hereby amended to include the following areas among those areas determined to have been adversely affected by the catastrophe declared a major disaster by the President in his declaration of December 4, 2002:

Kodiak Island Borough for Public Assistance (already designated for Individual Assistance).

Alaska Railroad right-of-way between Milepost 79 and Milepost 102 along the Turnagain Arm and state highway Milepost 4 Power Creek Road highway in the Cordova area for Public Assistance.

(The following Catalog of Federal Domestic Assistance Numbers (CFDA) are to be used for reporting and drawing funds: 83.537, Community Disaster Loans; 83.538, Cora Brown Fund Program; 83.539, Crisis Counseling; 83.540, Disaster Legal Services Program; 83.541, Disaster Unemployment Assistance (DUA); 83.556, Fire Management Assistance; 83.558, Individual and Household Housing; 83.559, Individual and Household Disaster Housing Operations; 83.560 Individual and Household Program—Other Needs; 83.544, Public Assistance Grants; 83.548, Hazard Mitigation Grant Program.)

**Joe M. Allbaugh,**  
Director.

[FR Doc. 03–4723 Filed 2–27–03; 8:45 am]

BILLING CODE 6718–02–P